

Isospectral Theory of Euler Equations

Yanguang (Charles) Li

Department of Mathematics,
University of Missouri
Columbia, MO 65211
E-mail: cli@math.missouri.edu

Roman Shvidkoy

Department of Mathematics,
University of Missouri
Columbia, MO 65211
E-mail: shvidkoy@math.missouri.edu

February 1, 2008

Abstract

Isospectral problem of both 2D and 3D Euler equations of inviscid fluids, is investigated. Connections with the Clay problem are described. Spectral theorem of the Lax pair is studied.

PACS Codes: 47, 02.

MSC numbers: 35, 51.

Keywords: Isospectral Theory, Lax Pair, Euler Equation.

1. Introduction

This note is a continuation of the works [1] and [2]. It focus upon the isospectral property of the Lax pairs of both 2D and 3D Euler equations. It provides efforts towards the connection between isospectral theory and the Clay problem on Navier-Stokes equations.

2. 2D Euler Equation

The 2D Euler equation can be written in the vorticity form,

$$\partial_t \Omega + \{\Psi, \Omega\} = 0 , \quad (2.1)$$

where the bracket $\{ , \}$ is defined as

$$\{f, g\} = (\partial_x f)(\partial_y g) - (\partial_y f)(\partial_x g) ,$$

where Ψ is the stream function given by,

$$u = -\partial_y \Psi , \quad v = \partial_x \Psi ,$$

where u and v are respectively the velocity components along x and y directions, and the relation between vorticity Ω and stream function Ψ is,

$$\Omega = \partial_x v - \partial_y u = \Delta \Psi .$$

Theorem 2.1 (Li, [1]) *The Lax pair of the 2D Euler equation (2.1) is given as*

$$\begin{cases} L\varphi = \lambda\varphi , \\ \partial_t \varphi + A\varphi = 0 , \end{cases} \quad (2.2)$$

where

$$L\varphi = \{\Omega, \varphi\} , \quad A\varphi = \{\Psi, \varphi\} ,$$

and λ is an imaginary constant, and φ is a complex-valued function.

2.1. Isospectral Theory and Conservation Laws

Denote by H^s the Sobolev space $H^s(R^2)$ or $H^s(T^2)$, and $\|\cdot\|_s$ the H^s norm.

Theorem 2.2 *Let Ω be a solution to the 2D Euler equation (2.1), φ be a solution to the Lax pair (2.2) at (Ω, λ) , then*

$$I = \frac{\|\{\Omega, \varphi\}\|_s}{\|\varphi\|_s}$$

is conservation law, i.e., I is independent of t .

Proof: Take the H^s norm on both side of the first equation in the Lax pair (2.2), then

$$I = \frac{\|\{\Omega, \varphi\}\|_s}{\|\varphi\|_s} = |\lambda|.$$

By the isospectral property of the Lax pair (2.2), I is independent of t . $\square$

An interesting idea is to try to use the conservation law I to prove global well-posedness. In this 2D Euler case, the idea is not very important since the global well-posedness is already proved. The hope is to use this idea to prove the global well-posedness of 3D Euler equation.

Lemma 2.1 *If φ solves the Lax pair (2.2), then $f(\varphi)$ solves*

$$\partial_t f(\varphi) + \{\Psi, f(\varphi)\} = 0, \quad (2.3)$$

for any f smooth enough.

Proof: The proof is completed by direct verification. $\square$

Lemma 2.2 *If $\{\phi_j\}_{j=1,2,\dots}$ is a complete base of H^s , where ϕ_j 's are $f(\varphi)$'s at different values of λ ; then*

$$\Omega = \sum_{j=1}^{\infty} a_j \phi_j,$$

where a_j 's are complex constants.

Proof: Since Equation (2.3) is a linear equation, the claim of Lemma 2.1 implies the current lemma. $\square$

2.2. The Spectrum of L

Denote by f_τ the flow generated by the vector field $(\Omega_y, -\Omega_x)$ on T^2 .

Theorem 2.3 *Denote by $\sigma_X(L)$ the spectrum of the operator L (2.2) in the space X .*

1. *If $\Omega \not\equiv \text{constant}$, then $\sigma_{L^2(T^2)}(L) = i\mathbb{R}$.*
2. *Denote by Λ the collection of all Lyapunov exponents of the flow f_τ , then $\Lambda/\{0\} + i\mathbb{R} \subset \sigma_{H^1(T^2)}(L)$.*

For a proof of this theorem, see [3].

3. 3D Euler Equation

The 3D Euler equation can be written in vorticity form,

$$\partial_t \Omega + (u \cdot \nabla) \Omega - (\Omega \cdot \nabla) u = 0, \quad (3.1)$$

where $u = (u_1, u_2, u_3)$ is the velocity, $\Omega = (\Omega_1, \Omega_2, \Omega_3)$ is the vorticity, $\nabla = (\partial_x, \partial_y, \partial_z)$, $\Omega = \nabla \times u$, and $\nabla \cdot u = 0$. u can be represented by Ω for example through Biot-Savart law.

Theorem 3.1 (Childress, [4]) *The Lax pair of the 3D Euler equation (3.1) is given as*

$$\begin{cases} L\varphi = \lambda\varphi, \\ \partial_t \varphi + A\varphi = 0, \end{cases} \quad (3.2)$$

where

$$L\varphi = (\Omega \cdot \nabla)\varphi - (\varphi \cdot \nabla)\Omega, \quad A\varphi = (u \cdot \nabla)\varphi - (\varphi \cdot \nabla)u,$$

λ is a complex constant, and $\varphi = (\varphi_1, \varphi_2, \varphi_3)$ is a complex 3-vector valued function.

Theorem 3.2 (Montgomery-Smith, [5]) *Another Lax pair of the 3D Euler equation (3.1) is given as* ■

$$\begin{cases} L\phi = \lambda\phi, \\ \partial_t \phi + A\phi = 0, \end{cases} \quad (3.3)$$

where

$$L\phi = (\Omega \cdot \nabla)\phi, \quad A\phi = (u \cdot \nabla)\phi,$$

λ is a complex constant, and ϕ is a complex scalar-valued function.

3.1. Isospectral Theory and Conservation Laws

Denote by H^s the Sobolev space $H^s(R^3)$ or $H^s(T^3)$, and $\|\cdot\|_s$ the H^s norm.

Theorem 3.3 *Let Ω be a solution to the 3D Euler equation (3.1), ϕ be a solution to the Lax pair (3.3) at (Ω, λ) , then*

$$I = \frac{\|(\Omega \cdot \nabla)\phi\|_s}{\|\phi\|_s}$$

is conservation law, i.e., I is independent of t .

Proof: Take the H^s norm on both side of the first equation in the Lax pair (3.2), then

$$I = \frac{\|(\Omega \cdot \nabla)\phi\|_s}{\|\phi\|_s} = |\lambda|.$$

By the isospectral property of the Lax pair (3.2), I is independent of t . □

Remark 3.1 *Of course, the significance of the conservation laws comes from their potential in providing a priori bounds and establishing the global well-posedness of 3D Euler equation, hence, of 3D Navier-Stokes equation (one of the Clay problems).*

Lemma 3.1 *If $\{\varphi_j\}_{j=1,2,\dots}$ is a complete base of H^s , where φ_j 's solve the Lax pair (3.2) at different values of λ ; then*

$$\Omega = \sum_{j=1}^{\infty} a_j \varphi_j ,$$

where a_j 's are complex constants.

Proof: The proof is completed by comparing the second equation in the Lax pair (3.2) and the 3D Euler equation. $\square$

References

- [1] Y. Li. A Lax Pair for the Two Dimensional Euler Equation. *J. Math. Phys.*, 42, No.8:3552–3553, 2001.
- [2] Y. Li and A. Yurov. Lax Pairs and Darboux Transformations for Euler Equations. *Submitted, available at: <http://xxx.lanl.gov/abs/math.AP/0101214>, or <http://www.math.missouri.edu/~cli>*, 2000.
- [3] Y. Latushkin and R. Shvidkoy. Spectral Properties of Linearized 2D Euler Equation and Evolution Semigroup. *Preprint (available upon request)*, 2002.
- [4] S. Childress. *personal communication*, 2000.
- [5] S. Montgomery-Smith. *personal communication*, 2002.